

## Key Targets

1

To optimise colostrum production by the sow

2

To make sure that piglets suckle within 6 hours of birth



3

To ensure that every piglet gets a fair share of the colostrum

(minimum 100ml in first 16h)



# Colostrum:

## Food for life

Colostrum is secreted from the udder immediately after farrowing and within several hours its composition changes to that representing sow milk. Colostrum contains essential antibodies (immunoglobulins) for the health of the newborn piglet; antibodies are produced by the body to kill bacteria and viruses. The piglet is born with very few of the protecting antibodies necessary to thrive and relies on the sow's colostrum to obtain them in the defence against bacteria and viruses.

Colostrum is a rich source of highly digestible nutrients, which are critical to establishing the newly born piglet and studies indicate that colostrum contains natural growth factors for the normal development of vital life-sustaining organs (eg brain, heart, pancreas, liver and kidneys) and the immature gut.

### COLOSTRUM INTAKE BY THE NEWBORN PIGLET

- Colostrum intake by the newborn piglet is highly variable and may range from **200–450g** per piglet
- Intake depends not only on the piglet's ability to extract colostrum but also on the ability of the sow to produce enough for the whole litter
- Many factors can influence both colostrum production and consumption, eg sow parity and nutrition, litter size and piglet vigour at birth
- Piglets with very low birth weights and which lack vigour at birth may not consume sufficient colostrum for survival or for their longer-term health and performance.



Colostrum is a rich source of highly digestible nutrients, which are critical to establishing the newly born piglet

### WHAT HAPPENS IF PIGLETS CONSUME LITTLE OR NO COLOSTRUM?

- The amount and timing of colostrum consumption are both critical to the subsequent health, development, survival and lifetime performance of the newborn piglet
- Colostrum intake must be optimised as soon as possible after farrowing
- After **6** hours the gut begins to 'close' and after **24** hours the full benefits of the immunoglobulins are no longer available to the piglet
- During this critical window, the piglet is able to rapidly absorb the protective colostrum factors (immunoglobulins) which are essential for good health and performance
- Unless sufficient colostrum has been consumed, the piglet remains highly vulnerable to disease
- The minimum threshold requirement for colostrum is unknown; however a general guide is that piglets will consume **150–280g/kg** body weight soon after birth.



## MANAGEMENT GUIDELINES

### How can colostrum production be optimised?

Little is known about the factors affecting colostrum production but the rearing strategy for replacement gilts, vaccination policy, and the disease and hygiene status of the farm are likely to influence the immunological properties of colostrum.

Apply best sow management principles to ensure high colostrum and milk yields and litter productivity:

- Reduce stress before, during and after farrowing
- Feed diets appropriate for pregnancy and lactation (see the Stotfold sow feeding strategy)
- Ensure sows have unrestricted access to fresh drinking water at all times.

### How can each piglet get a fair share of the colostrum?

#### ● Supervise farrowing:

Supervision during and immediately after farrowing should focus on early establishment of piglets at the teats to ensure adequate intake of colostrum; remember the ability to absorb antibodies falls rapidly after **6** hours and is almost completely gone **24** hours after birth

#### ● Batch farrowing:

This can assist in the planning and implementation of stockworker supervision of sows and piglets at farrowing

#### ● Synchronised farrowing:

The use of synchronising agents needs careful consideration and should be discussed with your veterinarian

#### ● Cross-fostering:

To ensure colostrum sharing, cross-fostering should take place as soon as possible following birth, ideally across sows within **24** hours or sooner after farrowing. If this is not possible, colostrum sharing in large litters should be facilitated by shift (split) suckling

- If porcine colostrum is unavailable, bovine colostrum may be used as a substitute
- Consider milking off surplus colostrum from some sows, freezing and using for compromised pigs (do not defrost using a microwave!). However, on most commercial farms, milking sows is not a practical option.

For more information on establishing the newborn pig, see *Action for Productivity 14: Newborn management*.



Colostrum intake must be optimised as soon as possible after farrowing



Assisted suckling



Split-suckling

© Agriculture and Horticulture Development Board 2017. No part of this publication may be reproduced in any material form (including by photocopy or storage in any medium by electronic means) or any copy or adaptation stored, published or distributed (by physical, electronic or other means) without the prior permission in writing of the Agriculture and Horticulture Development Board, other than by reproduction in an unmodified form for the sole purpose of use as an information resource when the Agriculture and Horticulture Development Board is clearly acknowledged as the source, or in accordance with the provisions of the Copyright, Designs and Patents Act 1988. All rights reserved.

While the Agriculture and Horticulture Development Board seeks to ensure that the information contained within this document is accurate at the time of printing, no warranty is given in respect thereof and, to the maximum extent permitted by law, the Agriculture and Horticulture Development Board accepts no liability for loss, damage or injury howsoever caused (including that caused by negligence) or suffered directly or indirectly in relation to information and opinions contained in or omitted from this document.

AHDB Pork is part of the Agriculture and Horticulture Development Board.